

Summer Wave® Torenia



Flowers all season, yielding a much higher bloom count than others. With a lush, trailing habit and vigorous growth, they're perfect for shady beds or hanging baskets. Self-cleaning and well-branching, they require very little maintenance. SUMMER WAVE is a registered trademark of SUNTORY FLOWERS LIMITED.

General & Information

Height:	Spread:	Exposure:
8-10" (20-25cm)	20-36" (51-91cm)	Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch: Yes	Average Propagation Time (weeks):	Comments:
50, 72, 84, 105	No	6-9		3-4	Reduce mist as soon as possible.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.8-6.2	5,000-8,000	71-79°F (22-26° C)	59-64°F (15-18° C)	175-225	1-2	Generally not needed.	Whitefly, spider mites, thrips	Botrytis	Torenia is a warm-season crop that prefers night temperatures above 60°F (16°C). Pinch regularly to shape.

Crop Time


4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 5-7 weeks	1-3 ppp, 6-8 weeks	3-5 ppp, 8-11 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Summer Wave® Varieties



Summer Wave® Large Amethyst



Summer Wave® Large Blue



Summer Wave® Large Violet