## **Sunbeam Bidens**



High vigor, early-season color and cool growing options make Sunbeam a must-have for large basket and combination programs.

Gene	ral & Infor	mation								
Height:		Spread:	Exposure:							
10-14"	(25-36cm)	18-28" (46-71cm)	Sun							
Propa	gation Gu	iide								
Tray Sizes		Rooting		Average Days with		Pinch:	Average Propagation Time	Comments:		
(cells/tray):		Hormone:		Mist:		Yes	(weeks):	Reduce and eliminate mist as soon as		
32, 50, 72, 84, 105		No		6-9		3-4		possible.		
Finish	ning									
Soil	Light	Temperature	Tempe	rature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
рН:	Levels	Day:	Night:		(ppm N):	Pinche	s: Regulators (S) =	Fungus gnats	stem rot bra	Pinch for better branching and
5.8-	(fc):	59-76°F (15-24°C	53-61°F	(12-16°C)	175-225	1-3	spray / (D) = drench:			
6.2	5,000-						B-Nine(S) 2,500 ppm. Also			shape.
	9,000						responds to Sumagic(S) 3-5			
							ppm or Sumagic(D) 0.25			
							ppm.			

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts : 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-7 weeks 2-3 ppp, 6-8 weeks 4-5 ppp, 6-9 weeks

**NOTE:** Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other Bidens Varieties



Bidens Sunbeam