Wasabi Coleus

Ball Flora Plant.

Stunning colors and patterns for full sun and shade. Versatile, premium varieties are very late to never flowering. Low-maintenance, high-impact plants are perfect for large pot programs and landscaping.

Durable and rugged for a coleus. Wasabi's serrated chartreuse leaves hold their brilliant color without fading.

ral & Infor	mation								
t:	Spread:	Exposure:							
(46-91cm)	16-28" (41-71cm)	Shade, Sun							
gation G	uide								
Sizes	Rooting	Average Days	Pinch:	Average Propagation Time (weeks):		Comments: Avoid stretch. Transplant on time. A B-Nine spray (1,500-2,000 ppm)			
tray):	Hormone:	with Mist:	No						
50, 72, 84, 105 No		4-6		2-3		10-12 days after stick can help to reduce stem elongation.			
ing									
Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Plant Growth		Diseases:	Comments:
Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S	Regulators (S) = spray / (D) = drench:		None	Pinch early to shape. Scout plants regularly
(fc):	74-85°F (23-29°	59-70°F (15-21°	175-225	0-2	spray / (D) = 0				
6.2 4,000-	C)	C)			Is responsive to E	B-Nine(S)	thrips		for whitefly. Avoid crowding at all stages
10,000									
					• • • •				
						Sumagic			
					(S) 5-10 ppm.				
Time									
-inch (10-	to 13-cm) Pots, Q	uarts : 6-inch (15	-cm) Pots, Ga	llons: 10- to	12-inch (25- to 3	30-cm) Tu	bs or Bask	ets:	
1 ppp, 5-7 weeks 1-3 p			, 6-8 weeks 4-5 ppp, 8-11 we						
	t: (46-91cm) (gation Gi Sizes tray): 84, 105 ing Light Levels (fc): 4,000- 10,000 Time -inch (10-	46-91cm) 16-28" (41-71cm) igation Guide Sizes Rooting sizes Hormone: 84, 105 No ing Light Temperature Levels Day: (fc): 74-85°F (23-29° 4,000- C) 10,000 Time	Spread: Exposure: (46-91cm) 16-28" (41-71cm) Shade, Sun Igation Guide Sizes Rooting Average Days sizes Rooting Average Days tray): Hormone: with Mist: 84, 105 No 4-6 sing Light Temperature Levels Day: Night: (fc): 74-85°F (23-29° 59-70°F (15-21° 4,000- C) C) C) 10,000 Time	Spread: Exposure: (46-91cm) 16-28" (41-71cm) Shade, Sun (gation Guide Sizes Rooting Average Days Pinch: (sizes Rooting Average Days Pinch: (tray): Hormone: with Mist: No 84, 105 No 4-6 ing Light Temperature Temperature Levels Day: Night: (ppm N): (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 4,000- C) C) C) 175-225 10,000 Time	Spread: Exposure: (46-91cm) 16-28" (41-71cm) Shade, Sun (gation Guide	t: Spread: Exposure: 46-91cm) 16-28" (41-71cm) Shade, Sun gation Guide Sizes Rooting Average Days Pinch: Average Propagation C tray): Hormone: with Mist: No Time (weeks): A 84, 105 No 4-6 2-3 1 ing Light Temperature Temperature Fertilization No. of Plant Growth Levels Day: Night: (ppm N): Pinches: Regulators (5 (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 0-2 spray / (D) = 4,000- C) C) C) Is responsive to 2,500-3,500 ppm Cycocel(S) 1,000 ppm tank mix or 3 (S) 5-10 ppm. Time -inch (10- to 13-cm) Pots, Quarts : 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 5	t: Spread: Exposure: 46-91cm) 16-28" (41-71cm) Shade, Sun tragation Guide Sizes Rooting Average Days Pinch: Average Propagation Comments tray): Hormone: with Mist: No Time (weeks): Avoid stretch. 84, 105 No 4-6 2-3 10-12 days af ting Light Temperature Temperature Fertilization No. of Plant Growth Levels Day: Night: (ppm N): Pinches: Regulators (S) = (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 0-2 Spray / (D) = drench: 10,000 C) C) C) C) Is responsive to B-Nine(S) 2,500-3,500 ppm and Cyccoel(S) 1,000-1,500 ppm tank mix or Sumagic (S) 5-10 ppm. Time Time Time	Spread: Exposure: (46-91cm) 16-28" (41-71cm) Shade, Sun rgation Guide Sizes Rooting Average Days Pinch: Average Propagation Comments: sizes No 4-6 2-3 Avoid stretch. Transplant or sign Light Temperature Fertilization No. of Plant Growth Pests: Light Temperature Night: (ppm N): Pinches: Regulators (S) = Whitefly, aphids, thrips (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 0-2 Spray / (D) = drench: aphids, thrips 4,000- C) C) C) C) Is responsive to B-Nine(S) 2,500-3,500 ppm and Cycocel(S) 1,000-1,500 ppm tank mix or Sumagic (S) 5-10 ppm. Time	t: Spread: Exposure: t46-91cm) 16-28" (41-71cm) Shade, Sun rgation Guide

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Coleus Varieties



Coleus Wasabi