SunGlow™ Nemesia



Blooms from early Spring to Summer and offers great heat tolerance. Easy to finish and perfect for quart production. Fragrant, delicate flowers are showstoppers in the garden in early Spring through Summer. SUNGLOW is a U.S. trademark of Penhow Nurseries Ltd.

General & Information

Height: Spread: Exposure:

12-14" (30-36cm) 10-12" (25-30cm) Sun

_			
- 0	rong	nation	. Guide

Tray Sizes	Rooting	Average Days	Pinch:	Average Propagation	Comments:
(cells/tray):	Hormone:	with Mist:	Recommended	Time (weeks):	Better with 2 cuttings per cell. Liner can be pinched at 3
72, 84, 105, 128	No	6-9		3-4	weeks or when well rooted.

72, 84,	105, 128	b, 128 NO 6-9 3-4 Weeks of		i wiieli well i	oolea.				
Finishing									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth Regulators	Pests:	Diseases:	Comments:
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	(S) = spray / (D) =	Thrips,	Root rot,	
5.7-	(fc):	64-75°F (18-24°C)	45-55°F (7-13°C)	175-225	1-2	drench:	aphids	Botrytis	
6.2	4,000-					If necessary, B-Nine spray at			
	7,000					2,500-3,500 ppm can be			
						effective. Growing cool helps to			
						prevent stretch.			
Cron	Timo								

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts : 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-7 weeks 2 ppp, 7-8 weeks 3-4 ppp, 9-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other SunGlow™ Varieties







SunGlow™ Yellow Bicolor