## Voltage™ White Osteospermum



The best Summer-performing osteo is great for growing in all climates. Traditional daisy has white petals with a bright yellow center. Blooms a little later than Voltage Yellow, with a more compact habit.

Spread:	Exposure:							
14-20" (36-51cm)	Sun							
ıide								
Tray Sizes Rooting (cells/tray): Hormone:		ys Pinch:	Average	verage Propagation		Comments:		
		Optional	Time (weeks):		Maintain uniformity in mist and soil temperature. Do not allow			
Yes	9-12	3-4			cuttings to wilt. Pinch to increase branching.			
Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:	
Day:	Night:	(ppm N):	Pinches:	Regulators (S) =	Aphids,	Root and	After transplanting, allow	
59-76°F (15-24°	44-55°F (7-13°C)	225-300	0-1	spray / (D) =	whitefly,	stem rot,	plants to become established	
C)				drench:	thrips	Botrytis	for 1-2 weeks at 60-65°F (15-	
			N p ta C p	Is responsive to B- Nine(S) 2,500-4,000 ppm alone or as a tank mix(S) with Cycocel 750-1,000 ppm. Bonzi(S) 15-30 ppm is also effective.			18°C). Once plants are well established, begin growing a recommended cool temperatures. Pinch optional. After 4-6 weeks of cool conditions, plants can be forced in 6-8 weeks.	
				.,				
	Rooting Hormone: Yes  Temperature Day: 59-76°F (15-24°	Rooting Average Da with Mist:  Yes 9-12  Temperature Day: Night:  59-76°F (15-24° 44-55°F (7-13°C)	Rooting	Rooting	Rooting Hormone: with Mist: Optional Time (weeks): Yes 9-12 3-4  Temperature Day: Night: (ppm N): Pinches: Regulators (S) = 59-76°F (15-24° C) C)  Temperature Temperature Fertilization (ppm N): Pinches: Regulators (S) = Sp-76°F (15-24° C) Sp-76°F (7-13°C) 225-300 Sp-76°F	Rooting Average Days Pinch: Average Propagation Comments Maintain unifo cuttings to will  Temperature Day: Night: (ppm N): Pinches: Regulators (S) = Sp-76°F (15-24° C)  Sp-76°F (15-24° A4-55°F (7-13°C) 225-300  Temperature Day: Night: (ppm N): Pinches: Regulators (S) = Spray / (D) = drench: Is responsive to B-Nine(S) 2,500-4,000 ppm alone or as a tank mix(S) with Cycocel 750-1,000 ppm. Bonzi(S) 15-30	Rooting Average Days With Mist: Yes 9-12 Temperature Day: Night: Sp-76°F (15-24° C) C)  Rooting Average Days Pinch: Average Propagation Optional Time (weeks): Maintain uniformity in mist and so cuttings to wilt. Pinch to increase cuttings to wilt	

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

3-5 ppp, 8-11 weeks

1-2 ppp, 8-11 weeks

## Other Osteospermum Varieties



1 ppp, 8-11 weeks

Osteospermum Voltage™ White