

# Vogue™ Double Petunia



Earlier to flower with a superior mounded, compact habit.  
Better branching than other double petunias on the market.  
More flowers in the center creates more flower power in containers and hanging baskets!

Bicolor is unique to the Vogue series.

## General & Information

| Height:         | Spread:          | Exposure: |
|-----------------|------------------|-----------|
| 8-10" (20-25cm) | 14-24" (36-61cm) | Sun       |

## Propagation Guide

| Tray Sizes<br>(cells/tray): | Rooting<br>Hormone: | Average Days<br>with Mist: | Pinch:<br>No | Average Propagation<br>Time (weeks): | Comments:  |
|-----------------------------|---------------------|----------------------------|--------------|--------------------------------------|--|
| 50, 72, 84, 105             | No                  | 6-9                        |              | 3-4                                  | Avoid stretch by moving crop to cooler air temperature in second and third weeks of propagation. Transplant on time. |

## Finishing

| Soil<br>pH: | Light<br>Levels<br>(fc): | Temperature<br>Day:   | Temperature<br>Night: | Fertilization<br>(ppm N): | No. of<br>Pinches: | Plant Growth<br>Regulators (S) =<br>spray / (D) =<br>drench:   | Pests:                       | Diseases:                                | Comments:   |
|-------------|--------------------------|-----------------------|-----------------------|---------------------------|--------------------|--|------------------------------|--|---|
| 5.4-<br>5.8 | 5,000-<br>8,000          | 59-76°F (15-24°<br>C) | 53-61°F (12-16°<br>C) | 225-300                   | 0-1                | B-Nine(S) 1,500-2,500 ppm, applied 2-4 times, can be used to control habit early; Bonzi(D) 0.25-1.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size. | Whitefly, thrips, leafminers | Botrytis, Root rot, Tobacco Mosaic Virus | Cool temperatures and high light will reduce or eliminate any need for PGRs. Monitor pH regularly for high pH and possible iron deficiency. Flowering will be best as days lengthen in late Spring and Summer. Daylength of more than 14 hours will hasten flowering. |

## Crop Time


| 4- to 5-inch (10- to 13-cm) Pots, Quarts : | 6-inch (15-cm) Pots, Gallons: | 10- to 12-inch (25- to 30-cm) Tubs or Baskets: |
|--|-------------------------------|--|
| 1 ppp, 6-8 weeks                           | 2-3 ppp, 7-9 weeks            | 3-5 ppp, 9-12 weeks                            |

**NOTE:** Growers should use the information presented here as guidelines only. Ball FloraPlant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball FloraPlant of any products listed herein. Ball FloraPlant's terms and conditions of sale shall apply to all products listed herein.

## Other Vogue™ Varieties




Vogue™ Blue




Vogue™ Lavender  
Vein



Vogue™ Lemon Berry



Vogue™ Pink



Vogue™ Red



Vogue™ White