Ball Flora Plant
Research & Development

725 Zenon Way
Arroyo Grande CA 93420
(805) 489-0228

Ipomoea Spotlight

Best Practices

Ball FloraPlant

Ipomoea Spotlight Direct Stick Best Practices

- Average Humidity: 65%
- Average moles/day: 16 M/m2d
- Soil pH: 5.7
- Fertilizer:

  17-5-17: 200 ppm N
- Container size:

  6-packs
- Stick week: 32
- Finish week: 36
- Total crop time: 4 weeks

Ball FloraPlant









Ball FloraPlant

2 Weeks After Transplant

Treatments:

- No Pinch/No PGR
- Pinch/No PGR: 2 weeks after stick
- Pinch/B-9 2,500 ppm: 2 weeks after stick
- No Pinch/B-9 2,500 ppm: 2 weeks after stick

Ball FloraPlant

2 Weeks After Stick

Control

Spotlight Black

Spotlight Red

Spotlight Lime

Ball FloraPlant

2 Weeks After Stick

Pinch / No PGR - 2 weeks after stick

Spotlight Black

Spotlight Red



Spotlight Lime

Ball FloraPlant

2 Weeks After Stick

No Pinch / B-Nine 2,500 ppm - 2 weeks after stick

Spotlight Black

Spotlight Red

Spotlight Lime

Ball FloraPlant

2 Weeks After Stick

Pinch / B-Nine 2,500 ppm - 2 weeks after stick

Spotlight Black



Spotlight Lime

Spotlight Red

Ball FloraPlant

4 Weeks After Transplant

Ball FloraPlant

4 Weeks After Stick

Control

Spotlight Black

Spotlight Red

Spotlight Lime

Ball FloraPlant

4 Weeks After Stick

Pinch / No PGR - 2 weeks after stick

Spotlight Black

Spotlight Red

Spotlight Lime

Ball FloraPlant

4 Weeks After Stick

No Pinch / B-Nine 2,500 ppm - 2 weeks after stick

Spotlight Black

Spotlight Red

Spotlight Lime

Ball FloraPlant

4 Weeks After Stick

Pinch / B-Nine 2,500 ppm - 2 weeks after stick

Spotlight Black

Spotlight Red

Spotlight Lime

Ball FloraPlant

Ipomoea Spotlight Best Practice

No Pinch/ No PGR

- Can be directly stuck into a quart
- Propagated directly in the greenhouse out of propagation
- No pinching or PGR's needed

Ball FloraPlant

Optional Treatments

Pinch/No PGR - 2 weeks after stick

No Pinch/B-Nine 2,500 – 2 weeks after stick

Pinch/B-Nine 2,500 – 2 weeks after stick

Ball FloraPlant