

Sunbeam Bidens



High vigor, early-season color and cool growing options make Sunbeam a must-have for large basket and combination programs.

General & Information

Height:	Spread:	Exposure:
10-14" (25-36cm)	18-28" (46-71cm)	Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks):	Comments:
32, 50, 72, 84, 105	No	6-9	Yes	(see note 1.) 3-4	Reduce and eliminate mist as soon as possible.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.8-6.2	5,000-9,000	59-76°F (15-24°C)	53-61°F (12-16°C)	175-225	1-3	B-Nine(S) 2,500 ppm. Also responds to Sumagic(S) 3-5 ppm or Sumagic(D) 0.25 ppm.	Fungus gnats	Root and stem rot	Pinch for better branching and shape.

Crop Time


4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 5-7 weeks	2-3 ppp, 6-8 weeks	4-5 ppp, 6-9 weeks

1. Propagation time:

Average propagation time for perennials includes 3-5 weeks to bulk-up the liner before beginning cold treatment when started in late Summer or early Autumn.

NOTE: Growers should use the information presented here as guidelines only. Ball FloraPlant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball FloraPlant of any products listed herein. Ball FloraPlant's terms and conditions of sale shall apply to all products listed herein.

Other Bidens Varieties



Bidens Sunbeam