

Campfire Coleus



Stunning colors and patterns for full sun and shade. Versatile, premium varieties are very late to never flowering. Low-maintenance, high-impact plants are perfect for large pot programs and landscaping.

Features eye-catching orange foliage and proven performance in the landscape, similar to Redhead and Wasabi. Stays vibrant in sun and shade. Late flowering, with a similar habit to Henna.

General & Information

| Height: | Spread: | Exposure: |
|------------------|------------------|------------|
| 14-28" (36-71cm) | 14-24" (36-61cm) | Shade, Sun |

Propagation Guide

| Tray Sizes (cells/tray): | Rooting Hormone: | Average Days with Mist: | Pinch: | Average Propagation Time (weeks): | Comments: |
|--------------------------|------------------|-------------------------|--------|-----------------------------------|--|
| 50, 72, 84, 105 | No | 4-6 | No | (see note 1.) 2-3 | Avoid stretch. Transplant on time. A B-Nine spray (1,500-2,000 ppm) 10-12 days after stick can help to reduce stem elongation. |

Finishing

| Soil pH: | Light Levels (fc): | Temperature Day: | Temperature Night: | Fertilization (ppm N): | No. of Pinches: | Plant Growth Regulators (S) = spray / (D) = drench: | Pests: | Diseases: | Comments: |
|----------|--------------------|--------------------|--------------------|------------------------|-----------------|---|--------------------------|-----------|--|
| 5.8-6.2 | 4,000-10,000 | 74-85°F (23-29° C) | 59-70°F (15-21° C) | 175-225 | 0-2 | Is responsive to B-Nine(S) 2,500-3,500 ppm and Cycocel(S) 1,000-1,500 ppm tank mix or Sumagic (S) 5-10 ppm. | Whitefly, aphids, thrips | None | Pinch early to shape. Scout plants regularly for whitefly. Avoid crowding at all stages. |

Crop Time

| 4- to 5-inch (10- to 13-cm) Pots, Quarts : | 6-inch (15-cm) Pots, Gallons: | 10- to 12-inch (25- to 30-cm) Tubs or Baskets: |
|--|-------------------------------|--|
| 1 ppp, 5-7 weeks | 1-3 ppp, 6-8 weeks | 4-5 ppp, 8-11 weeks |

1. Propagation time:

Average propagation time for perennials includes 3-5 weeks to bulk-up the liner before beginning cold treatment when started in late Summer or early Autumn.

NOTE: Growers should use the information presented here as guidelines only. Ball FloraPlant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball FloraPlant of any products listed herein. Ball FloraPlant's terms and conditions of sale shall apply to all products listed herein.

Other Coleus Varieties



Coleus Campfire