

# Celebrette New Guinea Impatiens



Includes 6 Grower's Choice varieties for the most uniform habit and timing across key colors.  
 Mid-size Celebrette delivers early, continuous flowering, with excellent branching, controlled habit and large blooms.  
 Thrives in both hot and cool conditions.  
 Ideal for small pot programs.

General & Information		
Height:	Spread:	Exposure:
8-10" (20-25cm)	10-12" (25-30cm)	Shade, Sun

Propagation Guide					
Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks): (see note 1)	Comments:
50, 72, 84, 105	No	7-8	No	3-4	Apply preventative fungicide spray after sticking. Reduce mist as soon as possible. Scout regularly for Botrytis.

Finishing									
Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.8-6.2	4,000-7,000	68-76°F (20-24°C)	59-64°F (15-18°C)	175-225	0	Bonzi(S) 2-10 ppm can be used to control habit early; Bonzi(D) 0.25-1.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size. Florel(S) 100-200 ppm can be used to improve branching but will usually not be needed.	Thrips, aphids, mites	Impatiens Necrotic Spot Virus, Botrytis, Pythium	Maintain 68°F (20°C) average daily temperature to optimize bloom size and reduce time to flower. Cooler temperatures will extend crop time. Grow crop somewhat dry to help control growth and hasten flowering. Avoid excess micro-nutrients in the soil. With good cultural practices and environmental conditions, no PGRs will be needed.

Crop Time		
4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 6-8 weeks	1-2 ppp, 7-9 weeks	4-5 ppp, 8-11 weeks


**1. Propagation time:**  
 Average propagation time for perennials includes 3-5 weeks to bulk-up the liner before beginning cold treatment when started in late Summer or early Autumn.

**NOTE:** Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other Celebrette Varieties



Celebrette Red Improved    Celebrette Bonfire Orange    Celebrette Lavender    Celebrette Strawberry Star    Celebrette Salmon Frost    Celebrette Hot Pink



Celebrette Frost    Celebrette Appleblossom