

Blazin' Rose Iresine



Striking multi-color leaves and strong vigor assure that the plant won't get lost in large mixed plantings; also ideal for landscapes.

General & Information

Height:	Spread:	Exposure:
18-30" (46-76cm)	16-20" (41-51cm)	Shade, Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks): (see note 1.)	Comments:
50, 72, 84, 105	No	4-7	No	3-4	Apply Bonzi spray 3-5 ppm on Blazin' Rose between Days 7 and 11 to control early stretch.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.8-6.2	4,000-7,000	71-79°F (22-26°C)	62-67°F (17-19°C)	175-225	1-2	Generally not needed. Responsive to Bonzi.	Whitefly	None	Pinch regularly to encourage branching and improve habit.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 7-9 weeks	2-3 ppp, 8-10 weeks	4-5 ppp, 10-13 weeks

1. Propagation time:

Average propagation time for perennials includes 3-5 weeks to bulk-up the liner before beginning cold treatment when started in late Summer or early Autumn.

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Iresine Varieties



Iresine Blazin' Rose