

Persian Shield Strobilanthes



Very vigorous with excellent heat tolerance. Works well in mixed containers, baskets and window boxes.

General & Information

Height:	Spread:	Exposure:
18-24" (46-61cm)	24" (61cm)	Partial Sun, Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch: Optional	Average Propagation Time (weeks): (see note 1.)	Comments:
50, 72, 84, 105	No	7-12		4-5	Pinch 18-21 days after sticking to improve branching. Allow root system to develop sufficiently before transplanting.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.8-6.2	3,000-6,000	74-79°F (23-26°C)	59-64°F (15-18°C)	175-225	1-3	Generally not needed.	Whitefly	Root rot	Moderate light levels produce more colorful foliage. Pinch early to shape. Feed regularly to maintain growth.

Crop Time


4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
	1-2 ppp, 8-10 weeks	3-5 ppp, 10-13 weeks

1. Propagation time:

Average propagation time for perennials includes 3-5 weeks to bulk-up the liner before beginning cold treatment when started in late Summer or early Autumn.

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Strobilanthes Varieties



Strobilanthes Persian Shield