

# Bumble Bee Blue Calibrachoa



Features a vibrant, unique bicolor pattern with large flowers and excellent heat performance.

Features a vibrant, unique bicolor pattern with large flowers and excellent heat performance. Habit matches Bumble Bee Pink.

## General & Information

**Height:** 10-15" (25-38cm)    **Spread:** 10-15" (25-38cm)    **Exposure:** Sun

## Propagation Guide

| Tray Sizes (cells/tray): | Rooting Hormone: | Average Days with Mist: | Pinch: Yes | Average Propagation Time (weeks): (see note 1.) | Comments:   |
|--------------------------|------------------|-------------------------|------------|---|---|
| 50, 72, 84, 105          | No               | 6-8                     |            | 3-4   | Avoid wide fluctuations in soil moisture. Apply preventative fungicide drench at 18-21 days after sticking. |

## Finishing

| Soil pH: | Light Levels (fc): | Temperature Day:  | Temperature Night: | Fertilization (ppm N): | No. of Pinches: | Plant Growth Regulators (S) = spray / (D) = drench:  | Pests:                     | Diseases: | Comments:  |
|----------|--------------------|-------------------|--------------------|------------------------|-----------------|--|----------------------------|-----------|--|
| 5.4-5.8  | 5,000-8,000        | 71-76°F (22-24°C) | 50-58°F (10-14°C)  | 225-300                | 1-2             | Use B-Nine(S) 2,500-5,000 ppm, Bonzi(D) 3-8 ppm or Florel(S) 300-500 ppm applied 1 time up to 3 times for large baskets to maintain habit as needed. | Thrips, aphids, leafminers | Root rot  | Cooler air temperatures and high light will reduce or eliminate any need for plant growth regulators. Maintain pH at 5.2-5.8 to optimize plant health. Crop times will be shortest as days lengthen in Spring. Apply preventative broad-spectrum fungicide drench early (7-10 days after transplant). Avoid saturated media and excessive feed levels. |

## Crop Time


**4- to 5-inch (10- to 13-cm) Pots, Quarts :** 1 ppp, 6-9 weeks    **6-inch (15-cm) Pots, Gallons:** 1-2 ppp, 7-11 weeks    **10- to 12-inch (25- to 30-cm) Tubs or Baskets:** 3-5 ppp, 8-12 weeks

### 1. Propagation time:

Average propagation time for perennials includes 3-5 weeks to bulk-up the liner before beginning cold treatment when started in late Summer or early Autumn.

**NOTE:** Growers should use the information presented here as guidelines only. Ball FloraPlant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball FloraPlant of any products listed herein. Ball FloraPlant's terms and conditions of sale shall apply to all products listed herein.

## Other Calibrachoa Varieties



Calibrachoa Bumble Bee Blue