AngelMist®



Big blooms have big impact in baskets and beds! Extreme heat and humidity are no match for these super-spreading plants.

Low growing for a unique groundcover.

Great blue bicolor addition to an already strong lineup. Summer's heat will transition the bicolor to a lavender blue two-tone.

| General & Information | | | | | | | |
|-----------------------|------------------|----------|--|--|--|--|--|
| Height: | Spread: | Exposure | | | | | |
| 4-10" (10-25cm) | 12-20" (30-51cm) | Sun | | | | | |

| Tray Sizes (cells/tray): 50, 72, 84, 105 | | Rooting | Avera | Average Days with | | Average Propagation Time (weeks): | | Comments: Avoid stretch. Pinch 18-21 days after | |
|--|--------|-----------------|-----------------|-------------------|----------|-----------------------------------|--------------|--|---|
| | | Hormone: Mist: | | | Yes | | | | |
| | | No | 7-9 | | | 3-4 | | sticking. | |
| Finish | ning | | | | | | | | |
| Soil | Light | Temperature | Temperature | Fertilization | No. of | Plant Growth | Pests: | Diseases: | Comments: |
| рН: | Levels | Day: | Night: | (ppm N): | Pinches: | Regulators (S) = | Whitefly, | Root rot, | High light will significantly |
| 5.8- | (fc): | 74-85°F (23-29° | 62-70°F (17-21° | 175-225 | 0-2 | spray / (D) = | caterpillars | Botrytis | improve plant quality by |
| 6.2 | 6,000- | C) | C) | | | drench: | | | encouraging branching, |
| | 10,000 | | | | | Generally not needed. | | | earlier flowering and the |
| | | | | | | Daminozide(S) 1,500- | | | ideal plant habit growth. |
| | | | | | | 3,000 ppm and | | | Best-suited to mid to late Spring and Summer |
| | | | | | | chlormequat chloride | | | programs. |
| | | | | | | 750-1,000 ppm tank mix | | | programo. |
| | | | | | | (S). Avoid ethephon use. | | | |

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 6-8 weeks 1-3 ppp, 7-9 weeks 3-5 ppp, 10-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other AngelMist® Varieties



Angelonia AngelMist® Spreading Pink



Angelonia AngelMist® Spreading Dark Purple



Angelonia AngelMist® Spreading Berry Sparkler



Angelonia AngelMist® Spreading Bluebird



Angelonia AngelMist® Spreading White