

- Thrives in extreme heat and humidity.
- Uniform habit and flowering time for easy, high-density production.
- Well-branched plants ship with less breakage, tangling and shrinkage.
- Works well in quarts, gallons and mixed containers. Also great in the landscape.

General & Information

Height: Spread: Exposure: 12-14" (30-36cm)10-12" (25-30cm)Sun

Propagation Guide

Tray Sizes (cells/tray):Rooting Hormone:Average Days with Mist:Pinch:Average Propagation Time (weeks):Comments:

50, 72, 84, 105 No 7-9 Yes 3-4 Avoid stretch. Pinch 18-21 days after sticking.

Florel use.

Finishing

Soil Light Temperature Temperature Fertilization No. of Plant Growth Regulators (S)Pests: Diseases: Comments: pH: Levels Night: Pinches: = spray / (D) = drench: Whitefly, Root rot, High light will significantly improve plant (ppm N): 5.8-(fc): 74-85°F (23-62-70°F (17-175-225 Generally not needed. Bcaterpillars Botrytis quality by encouraging branching, earlier 6.2 6,000- 29°C) 21°C) Nine(S) 1,500-3,000 ppm flowering and the ideal plant habit 10,000 and Cycocel 750-1,000 growth. Best suited to mid to late Spring ppm tank mix(S). Avoid and Summer programs.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :6-inch (15-cm) Pots, Gallons:10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-8 weeks 1-3 ppp, 7-9 weeks 3-5 ppp, 10-12 weeks

**NOTE:** Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other Archangel<sup>TM</sup> Varieties

