## **Burgundy Wedding Train**



Distinctive "wedding train" trailing habit. Foliage has dark color and unique texture. Great for shade containers and baskets. Hardy until frost and thrives in the heat.

|                  |                  | mation           |                 |               |   |  |   |                     |           |  |
|------------------|------------------|------------------|-----------------|---------------|---|--|---|---------------------|-----------|--|
| Height:          |                  | Spread:          | Exposure:       |               |   |  |   |                     |           |  |
| 10-12" (25-30cm) |                  | 24-36" (61-91cm) | Shade, Sun      |               |   |  |   |                     |           |  |
| Propa            | agation G        | uide             |                 |               |   |  |   |                     |           |  |
| Tray Sizes       |                  | Rooting          | Average Days    | Pinch:        | Average Propagation<br>Time (weeks):<br>2-3 |  | Comments:  Avoid stretch. Transplant on time. A daminozide spray (1,500-2,000 ppm) 10-12 days after stick can help to reduce stem elongation. |                     |           |  |
| (cells/tray):    |                  | Hormone:         | with Mist:      | No            |   |  |   |                     |           |  |
| 50, 72, 84, 105  |                  | No               | 4-6             |               |   |  |   |                     |           |  |
| Finish           | ning             |                  |                 |               |   |  |   |                     |           |  |
| Soil             | Light            | Temperature      | Temperature     | Fertilization | No. of                                      | Plant Growth   | Pes   | sts:                | Diseases: | Comments:  |
| H:               | Levels           | Day:             | Night:          | (ppm N):      | Pinches:                                    | Regulators (S) :   | = Thri  | ps,                 | Root rot  | Lava Rose Coleus will  |
| 5.8-             | (fc):            | 74-85°F (23-29°  | 59-70°F (15-21° | 175-225       | 0-2   | spray / (D) = dre  | ench: aph   | aphids,<br>whitefly |           | generally not require any<br>PGR applications during<br>production. Plants should<br>be pinched only as neede<br>since Lava Rose is<br>extremely free-branching. |
| 5.2              | 4,000-<br>10,000 | C)               | C)              |               |   | Generally not needed Uniconazole(S) 5-10 or a daminozide 2,50 3,500 ppm and chlormequat chloride -1,500 tank mix(S) or used to maintain half needed. | o) ppm<br>00-<br>e 1,000<br>can be  |                     |           |  |
| Crop             | Time             |                  |                 |               |   |  |   |                     |           |  |

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

3-4 ppp, 8-11 weeks

1-2 ppp, 6-8 weeks

## Other Coleus Varieties

1 ppp, 5-7 weeks



Coleus Burgundy Wedding Train