Florida Sun

General & Information

Height:	Spread:	Exposure:
12 - 18" (30 - 46 cm)	12 - 14" (30 - 36 cm)	Shade, Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks):	Comments:
50, 72, 84, 105	No	4-6	No	2-3	Avoid stretch. Transplant on time. A daminozide spray (1,500-2,000 ppm) 10-12 days after stick can help to reduce stem elongation.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.8-6.2	4,000-10,000	74-85°F (23-29° C)	59-70°F (15-21° C)	175-225	0-2	Generally not needed. uniconazole(S) 5-10 ppm or a daminozide 2,500-3,500 ppm and chlormequat chloride1,000-1,500 tank mix(S) can be used to maintain habit as needed.	Thrips, aphids, whitefly	Root rot	Lava Rose Coleus will generally not require any PGR applications during production. Plants should be pinched only as needed since Lava Rose is extremely free-branching.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 5-7 weeks	1-2 ppp, 6-8 weeks	3-4 ppp, 8-11 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Varieties



Coleus Florida Sun Rose