FlameThrower™



10- to 12-inch (25- to 30-cm) Tubs or Baskets:

Compact-to-medium vigor. Bold foliage colors and shapes. Add texture in small pots and mixed containers.

General & Information										
Heigh	t:	Spread:	Exposure:							
12-18"	(30-46cm)	16-18" (41-46cm)	Shade, Sun							
Propa	agation G	uide								
Tray Sizes		Rooting	Average Days	Pinch:	Average Propagation		Comments:			
(cells/tray):		Hormone:	with Mist:	No	Time (weeks):		Avoid stretch. Transplant on time. A daminozide spray (1,500-2,000			
50, 72, 84, 105		No	4-6		2-3		ppm) 10-12 days after stick can help to reduce stem elongation.			
Finish	ning									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Grow	th Regulators	Pests:	Diseases:	Comments:
рН:	Levels	Day:	Night:	(ppm N):	Pinches:	(S) = spray / (D) = drench:		Thrips, aphids, whitefly	Root rot	Scout plants regularly for whitefly. Avoid crowding at all stages.
5.8-	(fc):	74-85°F (23-29°	59-70°F (15-21°	175-225	0-2					
6.2	4,000-	C)	C)			Generally not needed. Uniconazole(S) 5-10 ppm or a				
	10,000									
							aminozide 2,500-3,500 ppm			
							uat chloride 1,000-			
						1,500 tank mix to maintain ha	x(S) can be used			
						to maintain na	ibit as needed.			

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

3-4 ppp, 8-11 weeks

6-inch (15-cm) Pots, Gallons:

1-2 ppp, 6-8 weeks

Other FlameThrower™ Varieties

4- to 5-inch (10- to 13-cm) Pots, Quarts :

Crop Time

1 ppp, 5-7 weeks

