FlameThrower™



10- to 12-inch (25- to 30-cm) Tubs or Baskets:

Compact-to-medium vigor. Bold foliage colors and shapes. Add texture in small pots and mixed containers.

Vibrant orange leaves with contrasting green edging.

Gene	ral & Infor	mation								
Heigh	t:	Spread:	Exposure:							
12-18" ((30-46cm)	16-18" (41-46cm)	Shade, Sun							
Propa	gation Gu	uide								
Tray Sizes (cells/tray): 50, 72, 84, 105		Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks): 2-3		Comments: Avoid stretch. Transplant on time. A daminozide spray (1,500-2,000 ppm) 10-12 days after stick can help to reduce stem elongation.			
				No						
Finish	ing									
Soil				Fertilization						
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Grov	th Regulators	Pests:	Diseases:	Comments:
	Light Levels	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Grov (S) = spray	•	Pests: Thrips,	Diseases: Root rot	Comments: Scout plants
рН:	•	•	•				•			
pH: 5.8-	Levels	Day:	Night:	(ppm N):	Pinches:	(S) = spray	/ (D) =	Thrips,		Scout plants regularly for whitefly. Avoid
pH: 5.8-	Levels (fc):	Day: 74-85°F (23-29°	Night: 59-70°F (15-21°	(ppm N):	Pinches:	(S) = spray drench: Generally not	/ (D) =	Thrips, aphids,		Scout plants regularly for whitefly. Avoid crowding at all
pH: 5.8-	Levels (fc): 4,000-	Day: 74-85°F (23-29°	Night: 59-70°F (15-21°	(ppm N):	Pinches:	(S) = spray drench: Generally not Uniconazole(staminozide 2,	needed. S) 5-10 ppm or a 500-3,500 ppm	Thrips, aphids,		Scout plants regularly for whitefly. Avoid
pH: 5.8-	Levels (fc): 4,000-	Day: 74-85°F (23-29°	Night: 59-70°F (15-21°	(ppm N):	Pinches:	(S) = spray drench: Generally not Uniconazole(staminozide 2) and chlormed	needed. needed. s) 5-10 ppm or a 500-3,500 ppm uat chloride 1,000-	Thrips, aphids,		Scout plants regularly for whitefly. Avoid crowding at all
Soil pH: 5.8- 6.2	Levels (fc): 4,000-	Day: 74-85°F (23-29°	Night: 59-70°F (15-21°	(ppm N):	Pinches:	(S) = spray drench: Generally not Uniconazole(staminozide 2, and chlormed 1,500 tank mi	needed. S) 5-10 ppm or a 500-3,500 ppm	Thrips, aphids,		Scout plants regularly for whitefly. Avoid crowding at all

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

3-4 ppp, 8-11 weeks

6-inch (15-cm) Pots, Gallons:

1-2 ppp, 6-8 weeks

Other FlameThrower™ Varieties

1 ppp, 5-7 weeks

4- to 5-inch (10- to 13-cm) Pots, Quarts :

