Limewire



Stunning colors and patterns for full sun and shade. Versatile, premium varieties are very late to never-flowering. Low-maintenance, high-impact plants are perfect for large pot programs and landscaping.

Deep burgundy ruffled leaves are edged in bright lime. Won't fade in full sun. Great branching.

General & Info	eneral & Information							
Height:	Spread:	Exposure						
14-28" (36-71cm)	14-28" (36-71cm)	Shade, Sun						

gation Gu	ide								
Sizes	Rooting	Average Days	Pinch: Average Propagation		agation Comment	Comments:			
(cells/tray): Hormone		with Mist:	No Time (weeks):		Avoid stretch	Avoid stretch. Transplant on time. A daminozide spray (1,500-2,000			
84, 105	No	4-6		2-3	ppm) 10-12 o	ppm) 10-12 days after stick can help to reduce stem elongation.			
ing									
Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:	
Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S) =	ench: aphids, ninozide thrips n and e(S)	None	Pinch early to shape. Scout plants regularly for whitefly. Avoid crowding at all stages.	
(fc):	74-85°F (23-29°	59-70°F (15-21°	175-225	0-2	spray / (D) = drench:				
4,000- 10,000	C)	C)			(S) 2,500-3,500 ppm and chlormequat chloride(S)				
	Sizes tray): 84, 105 sing Light Levels (fc): 4,000-	tray): Hormone: 84, 105 No iing Light Temperature Levels Day: (fc): 74-85°F (23-29° 4,000- C)	Sizes Rooting Average Days tray): Hormone: with Mist: 84, 105 No 4-6 sing Light Temperature Levels Day: Night: (fc): 74-85°F (23-29° 59-70°F (15-21° 4,000- C) C)	Sizes Rooting tray): Average Days with Mist: Pinch: No 84, 105 No 4-6 sing Temperature Fertilization Light Temperature Fertilization Levels Day: Night: (ppm N): (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 4,000- C)	Sizes Rooting tray): Average Days with Mist: Pinch: No Average Proportion 84, 105 No 4-6 2-3 sing Light Temperature Temperature Pertilization No. of Levels Day: Night: (ppm N): Pinches: (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 0-2 4,000- C) C) C) C) C)	Average Days	Average Days Pinch: Average Propagation Comments:	Rooting Average Days With Mist: No Time (weeks): Avoid stretch. Transplant on time. A daminozid ppm) 10-12 days after stick can help to reduce sing Light Temperature Temperature Fertilization Levels Day: Night: (ppm N): Pinches: Regulators (S) = Whitefly, None (fc): 74-85°F (23-29° 59-70°F (15-21° 175-225 0-2 spray / (D) = drench: aphids, 4,000- C) C) 10,000 C) C) C) Is responsive to daminozide (S) 2,500-3,500 ppm and chlormequat chloride(S)	


Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-7 weeks 1-3 ppp, 6-8 weeks 4-5 ppp, 8-11 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Coleus Varieties



Coleus Limewire