Henna



Stunning colors and patterns for full sun and shade. Versatile, premium varieties are very late to never flowering. Low-maintenance, high-impact plants are perfect for large pot programs and landscaping.

Very uniform. Serrated edges.

Leaves in chartreuse to copper have dark burgundy undersides.

Height:	al & Infor	Spread:	Exposure:							
•		•	•							
14-28" (3	6-/1cm)	14-16" (36-41cm)	Shade, Sun							
Propag	gation Gι	uide								
Tray Sizes		Rooting	Average Days	Pinch:	h: Average Propagation Time (weeks):		Comments: Avoid stretch. Transplant on time. A daminozide spray (1,500-2,000			
(cells/tray):		Hormone:	with Mist:	No						
50, 72, 84, 105		No	4-6		2-3		ppm) 10-12 days after stick can help to reduce stem elongation.			
Finishir	ng									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Grov	vth	Pests:	Diseases:	Comments:
рН:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators	Regulators (S) =		None	Pinch early to shape.
5.8-	(fc):	74-85°F (23-29°	59-70°F (15-21°	175-225	0-2	spray / (D) = drench:		Whitefly, aphids,		Scout plants regularly
6.2	4,000- 10,000	C)	C)			Is responsive	to daminozide	thrips		for whitefly. Avoid crowding at all stages.
						(S) 2,500-3,50	00 ppm and			
						chlormequat of	chlorideS)			
						1,000-1,500 p	pm tank mix or			
						uniconazole(S	S) 5-10 ppm.			

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

4-5 ppp, 8-11 weeks

1-3 ppp, 6-8 weeks

Other Coleus Varieties

1 ppp, 5-7 weeks



Coleus Henna