Honeybells

Part of the Bells Cuphea collection. Roots faster than comparable cuphea, reducing costs and production time. It can be produced at lower temperatures, decreasing the overall cost of Spring greenhouse production. Requires less special treatment because of its high resistance to Oedema.

Bursts of vibrant, hot pink flowers attract an abundance of pollinators, creating a thriving, lively garden. Its spreading habit makes it perfect for garden beds and stunning combinations. Honeybells will have your garden buzzing with beauty! A 2019 University of Georgia Plants of Distinction award winner.



| General & Info | eneral & Information | | | | | | |
|------------------|----------------------|------------------|--|--|--|--|--|
| Height: | Spread: | Exposure: | | | | | |
| 10-12" (25-30cm) | 14-20" (36-51cm) | Partial Sun, Sun | | | | | |

| 10-12" | (25-30cm) | 14-20" (36-51cm) | Partial Sun, Sun | | | | | | | |
|--|------------|---------------------|------------------|------------------------|---------------|--|---|-----------|-------------------------------------|--|
| Propa | agation Gu | uide | | | | | | | | |
| Tray Sizes Rooting (cells/tray): Hormone: 50, 72, 84, 105 No | | Average Days Pinch: | | Average Propagation Co | | Comments: | | | | |
| | | Hormone: | with Mist: | Yes | Time (weeks): | | Requires all flower buds to be pinched out of the liner in order to | | | |
| | | 6-8 | | 3-4 | | create better branching in finished product. | | | | |
| Finish | ning | | | | | | | | | |
| Soil | Light | Temperature | Temperature | Fertilization | No. of | Plant Growth | Pests: | Diseases: | Comments: | |
| pH: | Levels | Day: | Night: | (ppm N): | Pinches: | Regulators (S) = | Whitefly, | | Pinch as needed to shape. Fertilize | |
| 5.4- | (fc): | 70-75°F (21-24° | 59-64°F (15-18° | 175-225 | 1-2 | spray / (D) = drend | :h: aphids | | | |
| 5.8 6,000- | | C) | C) | | | Benefits from a paclobu | ıtrazol | | regularly. Crop times | |
| | 12,000 | | | | | (D) 2 ppm to control finis | | | will be shortest under | |
| | | | | | | habit. Also can benefit t | from a | | high light. | |
| | | | | | | Florel(S) 250 ppm tank i | mix | | | |
| | | | | | | with Configure(S) 75 pp | om to | | | |
| | | | | | | increase branching. | | | | |

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 6-8 weeks 2-5 ppp, 7-9 weeks 3-6 ppp, 10-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Cuphea Varieties



Cuphea Honeybells