

Cherrybells



Roots faster than comparable cuphea.
 Cuts costs and production time.
 Produce at lower temperatures.
 Spring greenhouse production costs less.
 Natural high resistance to Oedema.

General & Information

Height:	Spread:	Exposure:
10-12" (25-30cm)	14-20" (36-51cm)	Partial Sun, Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch: Yes	Average Propagation Time (weeks):	Comments:
50, 72, 84, 105	No	6-8		3-4	Requires all flower buds to be pinched out of the liner in order to create better branching in finished product.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.4-5.8	6,000-12,000	70-75°F (21-24° C)	59-64°F (15-18° C)	175-225	1-2	Benefits from a paclobutrazol (D) 2 ppm to control finished habit. Also can benefit from a ethephon(S) 250 ppm tank mix with benzyladenine(S) 75 ppm to increase branching.	Whitefly, aphids	Botrytis, Root rot	Pinch as needed to shape. Fertilize regularly. Crop times will be shortest under high light.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 6-8 weeks	2-5 ppp, 7-9 weeks	3-6 ppp, 10-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Cuphea Varieties



Cuphea Cherrybells