Tropica™ Modesto



Earlier blooming, better branching and more blooming capacity versus current market competitors.

Extreme heat, drought and weather tolerance.

Support not required.

General & Information Spread:

Exposure:

Propagation Guid Tray Sizes (cells/tray): 72-102		Rooting	Averaç	Average Days with Mist:		Average Propagation	Time	ime Comments:	
		Hormone:	Mist:			(weeks): 4-5		1 pinch in propagation, 1 in finish stage	
		Yes	12-14						
Finish	ning								
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S) =	Aphids (during	Fusarium,	
5.5-	(fc):	70-75°F (21-24°	60-70°F (16-21°C)	190-210	1	spray / (D) =	early Spring),	Pythium	
7.5	3,700-	C)				drench:	spider mites (durir	•	
	3.750					None	very dry weather)	

None

Crop Time 4- to 5-inch (10- to 13-cm) Pots, Quarts : 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 20-22 weeks 3 ppp, 20-22 weeks 1 ppp, 16-18 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Tropica™ Modesto Varieties



Dipladenia Tropica™ Modesto Pink



Dipladenia Tropica™ Modesto Red



Dipladenia Tropica™ Modesto White