

Tropica™ Grande



Earlier blooming, with better branching and blooming capacity versus the competition.

Ideal for patio or garden containers with a support (like a trellis or tepee).

Support required.

The world's first and only yellow dipladenia.
Hand-training is no longer required.

General & Information

Height:	Spread:	Exposure:
12-60" (30-152cm)	12-16" (30-41cm)	Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks):	Comments:
72-102	Yes	12-18	Yes	4-5	1 pinch in propagation, 1 in finish stage.

Finishing

Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.5-6.0	5,000-7,000	70-75°F (21-24° C)	60-65°F (16-18° C)	190-210	2	Daminozide(S) 2,500-5,000 ppm, paclobutrazol(D) 0.5-2 ppm for internode control. Benzyladenine(S) 300 ppm a week before pinch promotes branching.	Aphids (during early Spring), spider mites (during very dry weather)	Bacteria Leaf Spot	

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 16-18 weeks	1 ppp, 20-22 weeks	2 ppp, 20-22 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Tropica™ Grande Varieties



Dipladenia Tropica™ Grande Citrine