## **Marine**



Upright habit. Ideal for landscapes, baskets and patio planters.

Gene	ral & Infor	mation								
Heigh	t:	Spread:	Exposure:							
10-14"	(25-36cm)	12-18" (30-46cm)	Partial Sun							
Propa	agation Gu	uide								
Tray Sizes		Rooting	Average Days	Pinch:	Average Propagation Cor		comments:	mments:		
(cells/tray):		Hormone:	with Mist:	No	Time (weeks):		Stick immediately upon arrival. Apply preventative fungicide spray			
50, 72, 84, 105		Optional	6-9		3-4		after sticking. Reduce mist as soon as possible.			
Finish	ning									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:	
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S)	<ul> <li>Whitefly,</li> </ul>	Root rot	Cool temperatures and	
5.8-	(fc):	71-76°F (22-24°	50-61°F (10-16°	175-225	1-2	spray / (D) =	mites,		high light will reduce or	
6.2	5,000-	C)	C)			drench:	aphids		eliminate any need for plant growth regulators	
	9,000					Use daminozide 1,5	500-			
						3,000 ppm and				
						chlormequat chloric				
						750-1,000 ppm tanl	k mix			

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 6-7 weeks 3-5 ppp, 10-12 weeks 1-2 ppp, 8-9 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other Heliotrope Varieties



Heliotrope Marine