Marine



Upright plant performs well in landscapes, baskets and patio planters.

Gene	ral & Infor	mation								
Heigh	t:	Spread:	Exposure:							
10-14"	(25-36cm)	12-18" (30-46cm)	Partial Sun							
Propa	gation G	uide								
Tray Sizes		Rooting	Average Days	Pinch:	Average Pro	pagation	Comments:			
(cells/tray):		Hormone:	with Mist:	No	Time (weeks): 3-4		Stick immediately upon arrival. Apply preventative fungicide spray after sticking. Reduce mist as soon as possible.			
50, 72, 84, 105		Optional	6-9							
Finish	ing									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth Regulators (S) = spray / (D) =		Pests:	Diseases:	Comments:
рН:	Levels	Day:	Night:	(ppm N):	Pinches:			Whitefly, mites,	Root rot	Cool temperatures and high light will reduce or
5.8-	(fc):	71-76°F (22-24°	50-61°F (10-16°	175-225	1-2					
6.2	5,000-	C)	C)			drench:		aphids		eliminate any need for
	9,000					Use B-Nine 1,	500-3,000			plant growth regulators
						ppm and chlormequat				
						chloride 750-1 tank mix(S).	,000 ppm			
Cron	Tima					tarik mix(3).				
Crop		4- 42\ D-4- O	Augusta - Cimale (45	ama) Data Call	10 to 1	0 in ala (05 ta	20\ 7	Fulha au Daa	.l.ata.	
	•	to 13-cm) Pots, Q	•	cm) Pots, Gall		2-inch (25- to	3U-CM)	lubs of Bas	skets:	
1 ppp, 6-7 weeks 1-2				3-5 weeks 3-5		pp, 10-12 weeks				

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Heliotrope Varieties



Heliotrope Marine