Glimmer™

Ball Flora Plant...

This series with beautiful rose-like blooms replaces the Fiesta™ series with a double impatiens that is highly resistant to *Plasmopara destructor*, the cause of Impatiens downy mildew!

Although following a routine preventative spray treatment for downy mildew is still recommended, growers and consumers can be confident that these impatiens will continue to thrive and survive. Outstanding color item for baskets in the shade.

Vibrant pink and white double blooms offer high resistance to Impatiens downy mildew.

General & Information										
Height:		Spread:	Exposure:							
10-16" (25-41cm)		10-12" (25-30cm)	Shade, Partial Sun							
Propa	gation G	uide								
Tray Sizes		Rooting	Average Days with		Pinch:	inch: Average Propagation Time		Comments:		
(cells/tray):		Hormone:	: Mist:		No (weeks):		Reduce mist as soon as possible to reduce stretch.			
50, 72, 84, 105		No	5-8		2-3		Transplant on time.			
Finish	ning									
Soil	Light	Temperature	Temperature	Fertilization	No. o	Plant Growth	Pests:	Diseases:	Comments:	
pH:	Levels	Day:	Night:	(ppm N):	Pinch	es: Regulators (S) =	Leafminers, thrips	Impatiens Necrotic Spot Virus, Botrytis, Pythium	Start feeding immediately after transplanting. Avoid water stress in all stages of crop. Avoid cold stress (night temperatures below 65° F/18°C). Maintain EC above 1.0 (SME). Use early PGR applications to achieve ideal finished plants.	
5.8-	(fc): 4,000- 6,000	68-76°F (20-24° C)	56-61°F (13-16°	175-225	0-1	spray / (D) = drench:				
6.2			C)			Paclobutrazol(S) 5-10 ppm can be used to control habit early; paclobutrazol (D) 0.5-1.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size. Florel(S) 100-200 ppm can be used to improve branching but will usually not be needed.				
Crop	Time									
I- to 5	inch (10-	to 13-cm) Pots, Q	uarts : 6-inch	(15-cm) Pots, (Gallons:	10- to 12-inch (25- to 30-cm)	Tubs or Bas	kets:		

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

4-5 ppp, 9-12 weeks

1-2 ppp, 6-8 weeks

Other Glimmer™ Varieties

1 ppp, 5-7 weeks

