

Celebration



Vigorous and well branched.
Fills in fast with fewer plants per pot.
Full color range.
Excellent greenhouse performance.
Covered in large blooms, even in heat.

General & Information

Height: 10-12" (25-30cm)
Spread: 10-12" (25-30cm)
Exposure: Shade, Sun

Propagation Guide

| Tray Sizes (cells/tray): | Rooting Hormone: | Average Days with Mist: | Pinch: | Average Propagation Time (weeks): | Comments: |
|--------------------------|------------------|-------------------------|--------|-----------------------------------|---|
| 50, 72, 84, 105 | No | 7-8 | No | 3-4 | Apply preventative fungicide spray after sticking. Reduce mist as soon as possible. Scout regularly for Botrytis. |

Finishing


| Soil pH: | Light Levels | Temperature Day: | Temperature Night: | Fertilization (ppm N): | No. of Pinches: | Plant Growth Regulators (S) = spray / (D) = drench: | Pests: | Diseases: | Comments: |
|----------|-------------------|-------------------|--------------------|------------------------|-----------------|---|-----------------------|--|--|
| 5.8-6.2 | (fc): 4,000-7,000 | 68-76°F (20-24°C) | 59-64°F (15-18°C) | 175-225 | 0 | Paclobutrazol(S) 2-15 ppm can be used to control habit early; paclobutrazol(D) 0.25-2.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size. Ethephon(S) 100-200 ppm can be used to improve branching but will usually not be needed. | Thrips, aphids, mites | Impatiens Necrotic Spot Virus, Botrytis, Pythium | Maintain 68°F (20°C) average daily temperature to optimize bloom size and reduce time to flower. Cooler temperatures will extend crop time. Grow crop somewhat dry to help control growth and hasten flowering. Avoid excess micro-nutrients in the soil. With good cultural practices and environmental conditions, no PGRs will be needed. |

Crop Time


| 4- to 5-inch (10- to 13-cm) Pots, Quarts : | 6-inch (15-cm) Pots, Gallons: | 10- to 12-inch (25- to 30-cm) Tubs or Baskets: |
|--|-------------------------------|--|
| 1 ppp, 7-9 weeks | 1-2 ppp, 9-10 weeks | 3-5 ppp, 10-13 weeks |

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.


Other Celebration Varieties




New Guinea Impatiens
Celebration Icy Blue




New Guinea Impatiens
Celebration Apricot




New Guinea Impatiens
Celebration White




New Guinea Impatiens
Celebration Pink




New Guinea Impatiens
Celebration Bright
Salmon




New Guinea Impatiens
Celebration Purple Star




New Guinea Impatiens
Celebration Tropical
Peach




New Guinea Impatiens
Celebration Rose Star




New Guinea Impatiens
Celebration Electric
Rose



New Guinea Impatiens
Celebration Lavender
Glow



New Guinea Impatiens
Celebration Orange



New Guinea Impatiens
Celebration Deep Red