

- Bred to be extremely uniform in habit and flower timing across all colors, blooming all in the same week. Beneficial to greenhouse growers, as culture does not need adjusting. Since all colors finish together, they can be shipped to retail at the same time. Series truly performs like clockwork.
- · Ideal for quart and gallon containers, and hanging baskets.

Fresh color matches uniformity and timing.

General & Information

Height: Spread: Exposure: 8-10" (20-25cm)10-12" (25-30cm)Shade

Propagation Guide

Tray Sizes Rooting Average Days Pinch: Average Propagation Time Comments:

(cells/tray): Hormone: with Mist: No (weeks): Apply preventative fungicide spray after sticking. Reduce mist as soon as

50, 72, 84, 105 No 7-8 3-4 possible. Scout regularly for Botrytis.

Finishing

 $Soil Light \quad Temperature \ Temperature \ Fertilization \ No. \ of \quad Plant \ Growth \ Regulators \ (S) = \quad Pests: \quad Diseases: \quad Comments: \quad Pests: \quad Diseases: \quad Comments: \quad$

pH: Levels Day: Night: (ppm N): Pinches: spray / (D) = drench: Thrips, Impatiens Maintain 68°F (20°C) average daily

5.8 (fc): 68-76°F (20 59-64°F (15 175-225 0 Bonzi(S) 2-10 ppm can be used aphids, Necrotic temperature to optimize bloom size and

- 4,000--24°C) -18°C) to control habit early; Bonzi(D) mites Spot Virus, reduce time to flower. Cooler 6.2 7,000 Spot Virus, reduce time to flower. Cooler Botrytis, temperatures will extend crop time. Grow

matures will control final height Pythium crop somewhat dry to help control growth but not delay flowering or reduce flower size. Florel(S) 100-200 routrients in the soil. With good cultural

ppm can be used to improve practices and environmental conditions, no branching but will usually not be PGRs will be needed.

needed.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :6-inch (15-cm) Pots, Gallons:10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 6-8 weeks 1-2 ppp, 7-9 weeks 4-5 ppp, 8-11 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other ClockworkTM Varieties



New Guinea Impatiens ClockworkTM White ClockworkTM Orange ClockworkTM Orange Stripe

New Guinea Impatiens New Guinea Impatiens ClockworkTM Pink Star ClockworkTM Purple Stripe