## Blazin' Rose



Striking multicolor leaves and strong vigor assure that the plant won't get lost in large mixed plantings; also ideal for landscapes.

Gene	ral & Infor	mation							
Height:		Spread:	Exposure:						
18-30" (46-76cm)		16-20" (41-51cm)	Shade, Sun						
Propa	gation G	uide							
Tray Sizes (cells/tray):		Rooting Hormone:	Average Days with Mist:	Pinch: No	Average Propagation Time (weeks):		Comments: Apply paclobutrazol spray 3-5 ppm on Blazin' Rose between		
50, 72, 84, 105		No	4-7		3-4		Days 7-11 to control early stretch.		
Finish	ing								
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
<b>pH:</b> 5.8-6.2	Levels (fc): 4,000-	<b>Day:</b> 71-79°F (22-26°C	<b>Night:</b> C) 62-67°F (17-19°C)	(ppm N): 175-225	Pinches: 1-2	Regulators (S spray / (D) = drench:	) = Whitefly	None	Pinch regularly to encourage branching and improve habit.
	7,000					Generally not nee Responsive to paclobutrazol.	ded.		
Crop	Time								
4- to 5	-inch (10-	to 13-cm) Pots, Qu	uarts: 6-inch (15-c	m) Pots, Gallon	s: 10- to 12-ii	nch (25- to 30-cr	n) Tubs or Baske	ts:	
1 ppp, 7-9 weeks 2-3 ppp, 8-10 weeks					4-5 ppp, 10-1	13 weeks			

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other Iresine Varieties



Iresine Blazin' Rose