Blazin' Rose



drench:

Generally not needed. Responsive to paclobutrazol. and improve habit.

Striking multicolor leaves and strong vigor. Won't get lost in large mixed plantings. Ideal for landscapes.

General	&	Information

 Height:
 Spread:
 Exposure:

 18-30" (46-76cm)
 16-20" (41-51cm)
 Shade, Sun

_			_	
ם	ropad	nation	(211	แฝ๐

4,000-7.000

Tray Sizes		Rooting	Average Days	Pinch:	Average Propagation		Comments:		
(cells/	/tray):	Hormone:	with Mist:	No	Time (weeks):	: ,	Apply paclobutrazol:	spray 3-5 ppm o	n Blazin' Rose between
50, 72,	84, 105	No	4-7		3-4	[Days 7-11 to control	early stretch.	
Finishing									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S)	Whitefly	None	Pinch regularly to
5.8-	(fc):	71-79°F (22-26°C)	62-67°F (17-19°C)	175-225	1-2	spray / (D) =			encourage branching

Crop Time

6.2

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 7-9 weeks 2-3 ppp, 8-10 weeks 4-5 ppp, 10-13 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Iresine Varieties



Iresine Blazin' Rose