## $SunGlow^{TM} \\$



Blooms from early Spring to Summer and offers great heat tolerance. Easy to finish and perfect for quart production. Fragrant, delicate flowers are showstoppers in the garden in early Spring through Summer. SUNGLOW is a U.S. trademark of Penhow Nurseries Ltd.

General & Information

Height: Spread: Exposure: 12-14" (30-36cm)10-12" (25-30cm)Sun

Propagation Guide

Tray Sizes Rooting Average Days with Pinch: Average Propagation Time Comments:

(cells/tray): Hormone: Mist: Recommended(weeks): Better with 2 cuttings per cell. Liner can be pinched at 3

72, 84, 105, 128 No 6-9 3-4 weeks or when well rooted.

Finishing

Soil Light Temperature Temperature Fertilization No. of Plant Growth Regulators (S) = Pests: Diseases: Comments:

pH: Levels Night: (ppm N): Pinches: spray / (D) = drench: Thrips, Root rot, If grown cool and with high light 5.7- (fc): 64-75°F (18- 45-55°F (7- 175-225 1-2 If necessary, B-Nine spray at aphids Botrytis levels, SunGlow will generally require 6.2 4,000-2,500-3,500 ppm can be 24°C) 13°C) little, if any, PGRs. Avoid water-7,000 effective. Growing cool helps to logged media.

prevent stretch.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :6-inch (15-cm) Pots, Gallons:10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-7 weeks 2 ppp, 7-8 weeks 3-4 ppp, 9-12 weeks

**NOTE:** Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other SunGlow<sup>TM</sup> Varieties







Nemesia SunGlow<sup>TM</sup> Yellow Bicolor