## SunGlow™



spray at 2,500-3,500 ppm can be effective. Growing cool helps to prevent stretch. logged media.

Blooms from early Spring to Summer. Great heat tolerance. Easy to finish. Perfect for quart production. Fragrant, delicate showstopping flowers.

SUNGLOW is a U.S. trademark of Penhow Nurseries Ltd.

## General & Information

 Height:
 Spread:
 Exposure:

 12-14" (30-36cm)
 10-12" (25-30cm)
 Sun

Prop	agalion	Guide
T	C:	

•		Hormone:	Hormone: with Mist:	Recommended Time (weeks):  3-4		Better with 2 cuttings per cell. Liner can be pinched at 3 weeks or when well rooted.			
		No							
Finish	ning								
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S) =	Thrips,	Root rot,	If grown cool and with
5.7-	(fc):	64-75°F (18-24°	45-55°F (7-13°C)	175-225	1-2	spray / (D) =	aphids	Botrytis	high light levels, SunGlow
6.2	4.000-	C) .	, ,			drench:			will generally require little,
	7,000					If necessary, daminozide			if any, PGRs. Avoid water-

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-7 weeks 2 ppp, 7-8 weeks 3-4 ppp, 9-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Other SunGlow™ Varieties



Nemesia SunGlow™ Purple Bicolor



Nemesia SunGlow™ Yellow Bicolor