Bluerific

Unique color. Medium vigor. Well-suited for gallon and larger production.
Great addition to mixed containers. Early flowering.



General & Information

Height: Spread: Exposure:

8-10" (20-25cm) 16-22" (41-56cm)

Pro	nac	ıati	on	Gi	uida
110	vac	ıalı	UII	·Οι	มเนษ

Tray Sizes	Rooting	Average Days	Pinch: Average Propagation		Comments:
(cells/tray):	Hormone:	with Mist:	No	Time (weeks):	Avoid stretch by moving crop to cooler air temperature in second
50, 72, 84, 105	No	6-9		3-4	and third weeks of propagation. Transplant on time.

Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
pH: 5.4- 5.8	Levels (fc): 5,000- 8,000	Day: 59-76°F (15-24° C)	Night: 53-61°F (12-16° C)	(ppm N): 225-300	Pinches: 0-1	Regulators (S) = spray / (D) = drench: Daminozide(S) 1,000-2,000 ppm, applied 2-4 times, can be used to control habit early. Paclobutrazol(D) 0.25-1.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size.	Whitefly, thrips, leafminers	Botrytis, Root rot, Tobacco Mosaic Virus	Cool temperatures and high light will reduce need for PGRs. Monitor regularly for high pH and possible iron deficiency.

Crop Time

10- to 12-inch (25- to 30-cm) Tubs or Baskets: 4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons:

1 ppp, 5-7 weeks 2-3 ppp, 6-8 weeks 3-5 ppp, 8-11 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Petunia Varieties



Petunia Bluerific