Bluerific

Unique color.
Medium vigor.
Well-suited for gallon and larger production.
Great addition to mixed containers.
Early flowering.

General & Information

Height:	Spread:	Exposure:
8 - 10" (20 - 25 cm)	16 - 22" (41 - 56 cm)	Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Mist:	Pinch:	Average Propagation Time (weeks):	Comments:
50, 72, 84, 105	No	6-9	No	3-4	Avoid stretch by moving crop to cooler air temperature in second and third weeks of propagation. Transplant on time.

Finishing


Soil pH:	Light Levels (fc):	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases:	Comments:
5.4-5.8	5,000-8,000	59-76°F (15-24° C)	53-61°F (12-16° C)	225-300	0-1	Daminozide(S) 1,000-2,000 ppm, applied 2-4 times, can be used to control habit early. Paclobutrazol(D) 0.25-1.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size.	Whitefly, thrips, leafminers	Botrytis, Root rot, Tobacco Mosaic Virus	Cool temperatures and high light will reduce need for PGRs. Monitor regularly for high pH and possible iron deficiency.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :	6-inch (15-cm) Pots, Gallons:	10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 5-7 weeks	2-3 ppp, 6-8 weeks	3-5 ppp, 8-11 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Varieties



Petunia Bluerific