ColorRush™



Petunia ColorRush™

White

Cool temperatures and high

Notable for big vigor and excellent garden performance.

Our most vigorous line is ideal for 12-in. (30-cm) and larger containers and baskets, as well as landscapes.
The series will finish 10 to 12 weeks from stick, depending on region.

Night:

Masses of blooms hold up in the heat and rain.

Includes a range of bright, eye-catching colors and novelties.

These varieties have nicely matched habits, making production planning easier.

A 2019 University of Georgia Plants of Distinction award winner.

General & Information

Levels

Height: Spread: **Exposure:**

Day:

Propagation Guide Prop		-pu								
Tray Sizes Rooting Hormone: With Mist: Recommended 50, 72, 84, 105 No 6-9 Finishing Recommended Finishing Recommended Recommended 3-4 Recommen	10-12" (25-30cm)	24-36" (61-91cm)	Sun							
(cells/tray):Hormone:with Mist:RecommendedTime (weeks):Avoid stretch by moving crop to cooler air temperature in second and third weeks of propagation. Transplant on time.50, 72, 84, 105No6-93-4and third weeks of propagation. Transplant on time.	Propagation G	uide								
50, 72, 84, 105 No 6-9 3-4 and third weeks of propagation. Transplant on time. Finishing	Tray Sizes	Rooting	Average Days	Pinch:	Aver	age Propagation	Comments:			
Finishing	(cells/tray):	Hormone:	with Mist:	Recommended Tr		(weeks):	Avoid stretch by moving crop to cooler air temperature in second			
	50, 72, 84, 105	No	6-9	3-4		and third weeks of propagation. Transplant on time.				
Soil Light Temperature Temperature Fertilization No. of Plant Growth Pests: Diseases: Comments:	Finishing									
	Soil Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:	

(ppm N):

•		•	•	,		•	,,	2011/110, 11001	ooor tomporataroo ana mgi
5.4-	(fc):	59-76°F (15-24°	53-61°F (12-16°	225-300	0-1	spray / (D) =	thrips,	rot, Tobacco	light will reduce or eliminate
5.8	5.000-	C)	C)			drench:	leafminers	Mosaic Virus	any need for PGRs. Monitor
	8,000	,	,			Daminozide(S) 1,500- 2,500 ppm, applied 2-4 times, can be used to control habit early. Paclobutrazol(D) 0.25- 1.0 ppm applied as crop matures will control final height but not delay.			regularly for high pH and possible iron deficiency. Flowering will be best as days lengthen in late Spring and Summer. Daylength of more than 14 hours will hasten flowering.

Regulators (S) =

flowering or reduce flower size

Whitefly

Petunia ColorRush™

Watermelon Red

Botrytis Root

Pinches:

Crop Time

pH:

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 6-8 weeks 1 ppp, 7-9 weeks 3-5 ppp, 9-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other ColorRush™ Varieties

