Vogue™



Earlier to flower with a superior mounded, compact habit. Better branching than other double petunias on the market. More flowers in the center creates more flower power in containers and hanging baskets!

| Gene | ral & Info | rmation | | | | | | | |
|-----------------|-----------------|--------------------------------|----------------------------------|------------------------------|-----------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|--------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Heigh | nt: | Spread: | Exposure: | | | | | | |
| 8-10" (2 | 20-25cm) | 14-24" (36-61cm) | Sun | | | | | | |
| Propa | agation G | uide | | | | | | | |
| Tray Sizes | | Rooting | Average Da | ys Pinch: | Average Propagation Time (weeks): | | Comments: Avoid stretch by moving crop to cooler air temperature in second | | |
| (cells/tray): | | Hormone: | with Mist: | No | | | | | |
| 50, 72, 84, 105 | | No | 6-9 | | 3-4 | | and third weeks of propagation. Transplant on time. | | |
| Finish | hing | | | | | | | | |
| Soil | Light | Temperature | Temperature | Fertilization | No. of | Plant Growth | Pests: | Diseases: | Comments: |
| pH: | Levels | Day: 59-76°F (15-24° | Night: 53-61°F (12-16° | (ppm N): 225-300 | Pinches: 0-1 | Regulators (S) = spray / (D) = drench: | Whitefly, | rot, Tobacco | Cool temperatures and high light will reduce or eliminate any need for PGRs. Monitor pH regularly for high pH and possible iron deficiency. Flowering will be best as days lengthen in late Spring and Summer. Daylength of more than 14 hours will hasten flowering. |
| 5.4- | (fc): | | | | | | thrips, | | |
| 5.8 | 5,000- 8,000 | C) | C) | | | | leafminers | | |
| | | | | | | Daminozide(S) 1,500 2,500 ppm, applied 2 times, can be used to control habit early. Paclobutrazol(D) 0.2: 1.0 ppm applied as c matures will control f height but not delay flowering or reduce flower size. | -4 5 5- rop | | |

1 ppp, 6-8 weeks 2-3 ppp, 7-9 weeks 3-5 ppp, 9-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Vogue™ Varieties

4- to 5-inch (10- to 13-cm) Pots, Quarts :

Crop Time



Double Petunia Vogue™ Blue



Double Petunia Vogue™ Lavender



6-inch (15-cm) Pots, Gallons:

Double Petunia Vogue™ Lemon Berry



10- to 12-inch (25- to 30-cm) Tubs or Baskets:

Double Petunia Vogue™ Pink



Double Petunia Vogue™ Red



Double Petunia Vogue™ White