

CannonBall™



Perfect for premium pots and 10-in. (25-cm) hanging baskets.
 Uniform habit and timing.
 Cold tolerance for early-Spring sales.
 Excellent for cold-finishing programs.
 Compact habit for pot-tight production.
 Early flowering.
 All colors finish within a week of each other.
 Good branching without pinch or PGRs for growers looking for lower maintenance petunias.

This variety is more weather resilient than the original, and the timing is one week earlier to better match the series.

General & Information

Height: 6-8" (15-20cm) **Spread:** 8-12" (20-30cm) **Exposure:** Sun

Propagation Guide

| Tray Sizes (cells/tray): | Rooting Hormone: | Average Days with Mist: | Pinch: | Average Propagation Time (weeks): | Comments: |
|--------------------------|------------------|-------------------------|--------|-----------------------------------|--|
| 50, 72, 84, 105 | No | 6-9 | No | 3-4 | Avoid stretch by moving crop to cooler air temperature in second and third weeks of propagation. Transplant on time. |

Finishing

| Soil pH: | Light Levels (fc): | Temperature Day: | Temperature Night: | Fertilization (ppm N): | No. of Pinches: | Plant Growth Regulators (S) = spray / (D) = drench: | Pests: | Diseases: | Comments: |
|----------|--------------------|--------------------|--------------------|------------------------|-----------------|---|------------------------------|--|--|
| 5.4-5.8 | 5,000-8,000 | 59-76°F (15-24° C) | 53-61°F (12-16° C) | 225-300 | 0-1 | Daminozide(S) 1,500-2,500 ppm, applied 2-4 times, can be used to control habit early. Paclobutrazol(D) 0.25-1.0 ppm applied as crop matures will control final height but not delay flowering or reduce flower size. Daminozide applications late in the crop time can cause instability in flower color. | Whitefly, thrips, leafminers | Botrytis, Root rot, Tobacco Mosaic Virus | Cool temperatures and high light will reduce or eliminate any need for PGRs. Monitor regularly for high pH and possible iron deficiency. Flowering will be best as days lengthen in late Spring and Summer. Daylength of more than 14 hours will hasten flowering. |

Crop Time


| 4- to 5-inch (10- to 13-cm) Pots, Quarts : | 6-inch (15-cm) Pots, Gallons : | 10- to 12-inch (25- to 30-cm) Tubs or Baskets: |
|--|--------------------------------|--|
| 1 ppp, 5-7 weeks | 2-3 ppp, 6-8 weeks | 3-5 ppp, 8-11 weeks |

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other CannonBall™ Varieties



New Petunia CannonBall™ Blue 27 Petunia CannonBall™ Burgundy 26 Petunia CannonBall™ Red 26 Petunia CannonBall™ Yellow Petunia CannonBall™ Burgundy Star Petunia CannonBall™ Coral



Petunia CannonBall™ Pink Petunia CannonBall™ Rose Vein Petunia CannonBall™ White