Variegata

Trailing and vigorous. White-edged foliage. Use in combos and baskets.



waterings. Pinch regularly

to improve habit.

Rhizoctonia

General & Information

 Height:
 Spread:
 Exposure:

 8-10" (20-25cm)
 12-18" (30-46cm)
 Shade, Sun

Propagation Guid	е
------------------	---

5,000-

8 000

Tray Sizes		Rooting	Average Days with		Pinch:	Average Propagation	erage Propagation Time		Comments:	
(cells/tray):		Hormone:	Mist:		Optional	(weeks):		Apply preventative fungicide spray after		
50, 72, 84, 105		No	6-10		3-4			sticking.		
Finishing										
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:	
pH:	Levels	Day:	Night:	(ppm N):	Pinches	s: Regulators (S) =	Whitefly	Botrytis,	Regularly allow the media	
5.8-	(fc):	71-79°F (22-26°	56-61°F (13-16°C)	175-225	1-2	spray / (D) =	•	Pythium,	to dry between	

drench:

Generally not needed.

Crop Time

6.2

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 6-8 weeks 1-3 ppp, 7-10 weeks 3-5 ppp, 9-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Plectranthus Varieties



Plectranthus Variegata