## Variegata



Trailing and vigorous, this white-edged foliage plant is useful in combinations and baskets.

General & Information											
Heigh	t:	Spread:	Exposure:								
8-10" (2	20-25cm)	12-18" (30-46cm)	Shade, Sun								
Propa	agation G	uide									
Tray Sizes (cells/tray): 50, 72, 84, 105		Rooting Hormone: No		Average Days with Mist: <sup>6-10</sup>		Pinch: Optional	Average Propagation Time (weeks): 3-4		<b>Comments:</b> Apply preventative fungicide spray after sticking.		
						Finish					ning
Soil	Light	Temperatur	e Tempe	rature	Fertilization	No. of		Plant Growth	Pests:	Diseases:	Comments:
pH:	Levels	Day: Night:		(ppm N):	Pinche	es:	: Regulators (S) =	Whitefly	Botrytis,	Regularly allow the media	
5.8-	(fc):	71-79°F (22-26	° 56-61°F	(13-16°C)	175-225	1-2		spray / (D) =		Pythium,	to dry between
6.2	5,000-	C)						drench:		Rhizoctonia	waterings. Pinch regularly
	8,000							Generally not needed.			to improve habit.
Crop	Time										
4- to 5	5-inch (10-	- to 13-cm) Pots,	Quarts : 6	-inch (15	-cm) Pots, Gallo	ons: 10-	to 12-	nch (25- to 30-cm)	Tubs or Ba	skets:	
1 ppp, 6-8 weeks 1-				-3 ppp, 7-1	3-5	3-5 ppp, 9-12 weeks					

**NOTE:** Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Plectranthus Varieties



Plectranthus Variegata