Pink Brilliance

Heat and drought tolerant. Low maintenance. Bright pink, two-tone blooms.
Great for Summer basket programs.



General & Information

Height: Spread: **Exposure:**

12-20" (30-51cm) 24-28" (61-71cm) Sun

Propagation Gu	

	Tray Sizes	Rooting	Average Days	Pinch:	Average Propagation Time	Comments:
	(cells/tray):	Hormone:	with Mist:	Yes	(weeks):	Reduce mist as soon as possible. Maintain warm soil
	50, 72, 84, 105	Yes	10-15		3-4	temperature to optimize rooting.
Γ	Finishing					

Finish	ning								
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S) =	Whitefly,	Pythium,	Scaevola prefers a well-
5.8-	(fc):	74-79°F (23-26°	59-64°F (15-18°	175-225	1-2	spray / (D) =	leafminers	Rhizoctonia	drained soil. Allow the media
6.2	5,000-	C)	C)			drench:			to dry between waterings.
	10,000					Generally not needed. Paclobutrazol(D) 1-3 ppm or paclobutrazol (S) 20-40 ppm will control growth.			Night temperatures lower than recommended will significantly slow the crop and lead to problems with root rot.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts : 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

1 ppp, 5-7 weeks 1-2 ppp, 7-8 weeks 3-5 ppp, 10-12 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Scaevola Varieties



Scaevola Pink Brilliance