Persian Shield



Very vigorous. Excellent heat tolerance. Use in mixed containers, baskets and window boxes.

General & Information

Height: Spread: Exposure:

Partial Sun, Sun 18-24" (46-61cm) 24" (61cm)

Propagat	ion Guide
----------	-----------

Tray Sizes	Rooting	Average Days	Pinch:	Average Propagation	Comments:
(cells/tray):	Hormone:	with Mist:	Optional	Time (weeks):	Pinch 18-21 days after sticking to improve branching. Allow root
50, 72, 84, 105	No	7-12		4-5	system to develop sufficiently before transplanting.
Entra Lineau					

Finishing

Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:
pH:	Levels	Day:	Night:	(ppm N):	Pinches:	Regulators (S) =	Whitefly	Root rot	Moderate light levels produce
5.8-	(fc):	74-79°F (23-26°	59-64°F (15-18°C)	175-225	1-3	spray / (D) =			more colorful foliage. Pinch
6.2	3,000-	C)				drench:			early to shape. Feed
	6,000					Generally not needed.			regularly to maintain growth.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts: 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

> 1-2 ppp, 8-10 weeks 3-5 ppp, 10-13 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Strobilanthes Varieties



Strobilanthes Persian Shield