Persian Shield



Very vigorous with excellent heat tolerance. Works well in mixed containers, baskets and window boxes.

Gene	ral & Infor	mation									
Height:		Spread:	Exposure:								
18-24" ((46-61cm)	24" (61cm)	Partial Sun	, Sun							
Propa	gation G	uide									
Tray Sizes		Rooting		Average Days	Pinch:	Average Propagation		Comments:			
(cells/tray):		Hormone:		with Mist:	Optional Time (weeks):		F	Pinch 18-21 days after sticking to improve branching. Allow root			
50, 72, 84, 105		No		7-12	4-5		s	system to develop sufficiently before transplanting.			
Finish	ning										
Soil	Light	Temper	ature	Temperature -	Fertilization	No. of	Plant Growth	1	Pests:	Diseases:	Comments:
pH:	Levels	Day:	ı	Night:	(ppm N):	Pinches:	Regulators (S	S) =	Whitefly	Root rot	Moderate light levels produce more colorful foliage. Pinch early to shape. Feed regularly to maintain growth.
5.8-	(fc):	74-79°F (2	23-26° 5	59-64°F (15-18°C)	175-225	1-3	spray / (D) =				
6.2	3,000-	C)					drench:				
	6,000						Generally not ne	eded.			
Crop	Time										
		to 13-cm) P	ots Quart	s · 6-inch (15-	cm) Pots, Gall	ons: 10- to 12	inch (25- to 30)-cm) 1	Tubs or R	askets:	

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

3-5 ppp, 10-13 weeks

1-2 ppp, 8-10 weeks

Other Strobilanthes Varieties



Strobilanthes Persian Shield