

Wink



Excellent powdery mildew resistance – needs fewer chemical applications. Fast off the bench and onto retail shelves. Performs beautifully in baskets, mono and mixed containers.

General & Information

Height: Spread: Exposure:
8-10" (20-25cm) 12-18" (30-46cm) Sun

Propagation Guide

Tray Sizes (cells/tray):	Rooting Hormone:	Average Days with Pinch: Mist:	Average Propagation Time Yes (weeks):	Comments:
50, 72, 84, 105	No	6-9	3-4	Reduce and then eliminate mist as soon as possible. Apply preventative fungicide spray after sticking.

Finishing


Soil Light pH: Levels	Temperature Day:	Temperature Night:	Fertilization (ppm N):	No. of Pinches:	Plant Growth Regulators (S) = spray / (D) = drench:	Pests:	Diseases: Comments:
5.8- (fc): 6.2 5,000- 8,000	71-79°F (22- 26°C)	62-64°F (17- 18°C)	225-300	1-2	Responsive to a B-Nine 2,500-3,500 ppm/chlormequat chloride 750-1,000 ppm tank mix(S) and Florel(S) 250-300 ppm to control growth and improve plant habit.	Whitefly, leafminers, aphids	Root rot Regularly allow plants to dry between waterings. Pinch as needed to shape.

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:
1 ppp, 5-7 weeks 1-3 ppp, 7-8 weeks 4-5 ppp, 10-13 weeks

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

Other Wink Varieties



Verbena Wink
Burgundy



Verbena Wink Violet