Cabaret[®]



Cabaret carries the best-performing lineup of solid colors and exciting bicolors with early flowering!

Most varieties flower with a 10.5-hour day length, and stay open and in flower under low light conditions — a big plus at retail.

All new genetics flower early when produced for target production weeks prior to Week 5 in the South and Week 14 in the North.

Works well in 4.5 to 6-in. (11 to 15-cm) pots up to large baskets.

Brighter color and earlier to flower than the current Rose, with early flowering, so it is great to target production for week 14 or earlier. Perfect for tight basket production.

General & Information

Height: Spread: Exposure:

6-10" (15-25cm) 10-12" (25-30cm) Sun

Tray Sizes (cells/tray): 50, 72, 84, 105		Rooting	Average D	ays Pinch	: Average	Average Propagation Time (weeks): 3-4		Comments: Avoid wide fluctuations in soil moisture. Apply preventative fungicide drench at 18-21 days after sticking.		
		Hormone:	with Mist:	Yes	Time (we					
Finish	ning									
Soil	Light	Temperature	Temperature	Fertilization	No. of	Plant Growth	Pests:	Diseases:	Comments:	
pH: 5.4-5.8	Levels (fc): 5,000- 8,000	Day: 71-76°F (22-24° C)	Night: 50-58°F (10-14° C)	(ppm N): 225-300	Pinches: 1-2	Regulators (S) = spray / (D) = drench: Use daminozide(S)	Thrips, aphids, leafminers	aphids,	Cooler air temperatures and high light will reduce or elimina any need for plant growth regulators. Maintain pH at 5.2-5.8 to optimize plant health. Crop times will be shortest as days lengthen in Spring. Apply preventative broad-spectrum fungicide drench early (7-10 days after transplant). Avoid saturated media and excessiv feed levels.	
						2,500-5,000 ppm, paclobutrazol(D) 3-8 ppm or ethephon(S) 300-500 ppm applied				
						1 time up to 3 times for large baskets to maintain habit as needed.				

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

3-5 ppp, 8-12 weeks

1-2 ppp, 7-11 weeks

Other Cabaret® Varieties

1 ppp, 6-9 weeks



