# Cabaret®

Best-performing lineup of solid colors and bicolors with early flowering.
Most flower with a 10.5-hour day length.
Stays open and in flower under low light.
Early flowering; target production weeks prior to Week 5 (South) and Week 14 (North).
Works well in 4.5 to 6-in. (11 to 15-cm) pots up to large baskets.

Features pink flowers with a vein and an attractive pattern.

## General & Information

| Height: | Spread: | Exposure: |
|---|---|---|
| 6 - 10" (15 - 25 cm) | 10 - 12" (25 - 30 cm) | Sun |

## Propagation Guide

| Tray Sizes (cells/tray): | Rooting Hormone: | Average Days with Mist: | Pinch: | Average Propagation Time (weeks): | Comments: |
|---|---|---|---|---|---|
| 50, 72, 84, 105 | No | 6-8 | Yes | 3-4 | Avoid wide fluctuations in soil moisture. Apply preventative fungicide drench at 18-21 days after sticking. |

## Finishing


| Soil pH: | Light Levels (fc): | Temperature Day: | Temperature Night: | Fertilization (ppm N): | No. of Pinches: | Plant Growth Regulators (S) = spray / (D) = drench: | Pests: | Diseases: | Comments: |
|---|---|---|---|---|---|---|---|---|---|
| 5.4-5.8 | 5,000-8,000 | 71-76°F (22-24° C) | 50-58°F (10-14° C) | 225-300 | 1-2 | Use daminozide(S) 2,500-5,000 ppm, paclobutrazol(D) 3-8 ppm or ethephon(S) 300-500 ppm applied 1 time up to 3 times for large baskets to maintain habit as needed. | Thrips, aphids, leafminers | Root rot | Cooler air temperatures and high light will reduce or eliminate any need for plant growth regulators. Maintain pH at 5.2-5.8 to optimize plant health. Crop times will be shortest as days lengthen in Spring. Apply preventative broad-spectrum fungicide drench early (7-10 days after transplant). Avoid saturated media and excessive feed levels. |

## Crop Time


| 4- to 5-inch (10- to 13-cm) Pots, Quarts : | 6-inch (15-cm) Pots, Gallons: | 10- to 12-inch (25- to 30-cm) Tubs or Baskets: |
|---|---|---|
| 1 ppp, 6-9 weeks | 1-2 ppp, 7-11 weeks | 3-5 ppp, 8-12 weeks |


**NOTE:** Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

## Varieties



Calibrachoa Cabaret® Strawberry Parfait


Calibrachoa Cabaret® Early Yellow


Calibrachoa Cabaret® Classic White


Calibrachoa Cabaret® Early Hot Pink


Calibrachoa Cabaret® Early Neon Rose


Calibrachoa Cabaret® Classic Deep Blue


**New** Calibrachoa Cabaret® Early Orange Kiss



Calibrachoa Cabaret® Early Lemon Yellow


Calibrachoa Cabaret® Classic Bright Red


Calibrachoa Cabaret® Early Midnight Blue


Calibrachoa Cabaret® Early Light Pink



Calibrachoa Cabaret® Early Pink Star




Calibrachoa Cabaret® Early Strawberry Parfait




**New** Calibrachoa Cabaret® Early Frosty Lemon




Calibrachoa Cabaret® Early Orange




Calibrachoa Cabaret® Early Bright Orange



Calibrachoa Cabaret® Early Lavender



Calibrachoa Cabaret® Early Good Night Kiss




Calibrachoa Cabaret® Light Pink Kiss




Calibrachoa Cabaret® Early Red



Calibrachoa Cabaret® Classic Deep Yellow




Calibrachoa Cabaret® Classic Purple



Calibrachoa Cabaret® Classic Pink




Calibrachoa Cabaret® Early Lavender Kiss



Calibrachoa Cabaret® Early Midnight Kiss



Calibrachoa Cabaret® Early Bright White



Calibrachoa Cabaret® Diva Pink