## Conga™



Naturally compact habit is perfect for pot-tight production, with no pinching or PGRs.

Great choice for quarts, color bowls and window boxes.

Easy to grow - very uniform.

Excellent pH tolerance and profuse flowering give Conga superior hold at retail.

Flowers one week earlier than Cabaret Series.

Exciting bicolor is unique to the market, expanding the appeal of the series

General & Information Height: Spread: **Exposure:** 6-10" (15-25cm) 8-10" (20-25cm) Sun **Propagation Guide Tray Sizes** Rooting **Average Days** Pinch: **Average Propagation** Comments: with Mist: (cells/tray): Hormone: Time (weeks): Yes Avoid wide fluctuations in soil moisture. Apply preventative fungicide drench at 18-21 days after sticking. 50, 72, 84, 105 No 6-8 3-4 Finishing Soil Light Temperature Temperature **Fertilization** No. of **Plant Growth** Pests: Diseases: Comments: Pinches: Regulators (S) = pH: Levels Day: Night: (ppm N): Thrips. Root rot Cooler air temperatures and spray / (D) = (fc): aphids, high light will reduce or 5.4-71-76°F (22-24° 50-58°F (10-14° 225-300 1-2 drench: leafminers eliminate any need for plant 5.8 C) C) 5,000growth regulators. Maintain 8.000 Plant growth regulators pH at 5.2-5.8 to optimize are generally not needed plant health. Crop times will and should only be be shortest as days applied if plants show lengthen in Spring. Apply signs of requiring extra preventative broadcontrol. Daminozide(S) spectrum fungicide drench. 1,500-3,000 ppm applied 1 time up to 3 times should provide adequate control Crop Time 4- to 5-inch (10- to 13-cm) Pots, Quarts : 6-inch (15-cm) Pots, Gallons: 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

NOTE: Growers should use the information presented here as guidelines only. Ball Floraplant recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Ball Floraplant of any products listed herein. Ball Floraplant's terms and conditions of sale shall apply to all products listed herein.

1-3 ppp, 6-10 weeks

3-5 ppp, 8-12 weeks

## Other Conga™ Varieties

1 ppp, 5-8 weeks

